

THE IDEAL ENVELOPE OF AN OPERATOR ALGEBRA

DAVID P. BLECHER AND MASAYOSHI KANEDA

ABSTRACT. A left ideal of any C^* -algebra is an example of an operator algebra with a right contractive approximate identity (r.c.a.i.). Conversely, we show here and in [6] that operator algebras with r.c.a.i. should be studied in terms of a certain left ideal of a C^* -algebra. We study operator algebras and their multiplier algebras from the perspective of ‘Hamana theory’ and using the multiplier algebras introduced by the first author.

*This research was supported by a grant from the National Science Foundation.
Revision of Oct 8, 2001.

1. INTRODUCTION AND NOTATION

A left ideal of any C^* -algebra is an example of an operator algebra with a right contractive approximate identity. Conversely, we study operator algebras with right contractive approximate identity in terms of a certain left ideal of a C^* -algebra.

A (concrete) operator algebra is a subalgebra of $B(H)$, for some Hilbert space H . More abstractly, an operator algebra will be an algebra A with a norm defined on the space $M_n(A)$ of $n \times n$ matrices with entries in A , for each $n \in \mathbb{N}$, such that there exists a *completely isometric*¹ homomorphism $A \rightarrow B(H)$ for some Hilbert space H . In this paper all our operator algebras and spaces will be taken to be complete. We shall say that an operator algebra is *unital* if it has a two-sided contractive identity. In the present paper we are concerned with operator algebras with a one-sided (usually right) contractive approximate identity. We shall abbreviate ‘right (resp. left) contractive approximate identity’ to ‘r.c.a.i.’ (resp. ‘l.c.a.i.’).

In §2 of our paper we consider a certain ‘transference principle’, which can allow one to deduce many general results about operator algebras with r.c.a.i., from results about left ideals in a C^* -algebra (see the companion paper [6]). Namely there is an important left ideal $\mathfrak{J}_e(A)$ of a C^* -algebra $\mathcal{E}(A)$, which is associated to any such operator algebra. We call $\mathfrak{J}_e(A)$ the ‘left ideal envelope’ of A . This is analogous to the case of operator algebras with 2-sided c.a.i., which are largely studied these days in terms of a certain C^* -algebra, namely the C^* -envelope.

In §3 we study the ‘left multiplier operator algebra’ of an operator algebra A with r.c.a.i. (which will be a symmetrical theory to that of $RM(A)$ for an operator algebra A with l.c.a.i.). This theory does not work out quite as nicely as the case when A has a two-sided c.a.i., unless A is a left ideal of a C^* -algebra. We present some examples to show that some of the obvious candidate descriptions of this multiplier algebra are defective. A main motivation for studying multiplier algebras is because of their intimate connection with the extremely important notion of the ‘unitization’ of a nonunital algebra. A good ‘unitization’ procedure should reduce many problems about nonunital algebras to the well understood unital case. Although we are not at this point able to completely analyze the ‘unitization’ procedure for algebras with one-sided c.a.i., we at least do some of the groundwork here.

We end the introduction with some more notation, and some background results which will be useful in various places. We reserve the letters H, K for Hilbert spaces, and J for a left ideal of a C^* -algebra. If A is an algebra then we write $\lambda : A \rightarrow Lin(A)$ for the canonical ‘left regular representation’ of A on itself. If S is a subalgebra of A then the left idealizer of S is the subalgebra $\{x \in A : xS \subset S\}$ of A . Note S is a left ideal in this subalgebra, whence the name. Similarly for the right idealizer; the (2-sided) idealizer is the intersection of the left and right idealizer.

By a ‘representation’ $\pi : A \rightarrow B(H)$ of an operator algebra A we shall mean a completely contractive homomorphism. If A has r.c.a.i. and if we say that π is *nondegenerate*, then at the very least we mean that $\pi(A)H$ is dense in H . Note that this last condition does not imply in general that $\pi(e_\alpha)\zeta \rightarrow \zeta$ for $\zeta \in H$, where $\{e_\alpha\}$ is the r.c.a.i., as one is used to in the two-sided case.

Lemma 1.1. *Let A be an operator algebra with either a l.c.a.i. or a r.c.a.i.. Suppose that $\pi : A \rightarrow B(H)$ is a completely contractive representation, set K to be the norm closure of*

¹A map $T : X \rightarrow Y$ is completely isometric if $[x_{ij}] \mapsto [T(x_{ij})]$ is isometric on $M_n(X)$ for all $n \in \mathbb{N}$

$\pi(A)H$, and let $\pi' : A \rightarrow B(K)$ be the induced representation. Then π' is a completely contractive homomorphism, and it is nondegenerate in the sense that $\pi'(A)K$ is dense in K . Moreover if A has a r.c.a.i. and π is isometric (resp. completely isometric, 1-1), then π' is also isometric (resp. completely isometric, 1-1).

Consequently, if A is a subalgebra of $B(H)$ with a r.c.a.i., then we may view A as a nondegenerate subalgebra of $B(K)$, where K is the closure of AH .

Proof. That π' is a completely contractive homomorphism is clear. Clearly $\pi'(A)K \subset K$, however if $a \in A$ and $\zeta \in H$ then $\pi(a)\zeta = \lim \pi'(a)\pi(e_\alpha)\zeta \in \pi'(A)K$, if A has a r.c.a.i.. Taking linear combinations of such expressions $\pi(a)\zeta$, and limits, shows that any element of K is in the closure of $\pi'(A)K$. So π' is nondegenerate in this case. A similar argument holds if A has a l.c.a.i..

To see the statement here about the isometry note that for $a \in A$ we have

$$\|a\| = \|\pi(a)\| = \sup\{\|\pi(a)\zeta\| : \zeta \in \text{Ball}(H)\} = \sup \lim_{\alpha} \{\|\pi'(a)\pi(e_\alpha)\zeta\| : \zeta \in \text{Ball}(H)\}.$$

However the right hand side is dominated by $\|\pi'(a)\|$, so that π' is an isometry. A similar argument works for a complete isometry. The 1-1 assertion is easier. $\square$

If an operator algebra has only a l.c.a.i. then the ‘isometric’ assertions of the last result are not true in general. For a counterexample consider $A = R_n$, the subalgebra of M_n supported on the first row.

We will use without comment several very basic facts from C^* -algebra theory (see e.g. [19]), such as the basic definitions of the left multiplier algebra $LM(A)$, and the multiplier algebra $M(A)$, of a C^* -algebra. As a general reference for operator spaces the reader might consult [11, 20] or [17]. We write $\hat{} : X \rightarrow X^{**}$ for the canonical map, this is a complete isometry if X is an operator space, and is a homomorphism if X is an operator algebra, giving the second dual the Arens product [10].

We will often consider the basic examples C_n (resp. R_n) of operator algebras with right (resp. left) identity of norm 1; namely the $n \times n$ matrices ‘supported on’ the first column (resp. row). This is a left (resp. right) ideal of M_n , and has the projection E_{11} as the 1-sided identity.

If X and Y are subsets of an operator algebra we usually write XY for the *norm closure* of the set of finite sums of products xy of a term in X and a term in Y . For example, if J is a left ideal of a C^* -algebra A , then with this convention J^*J will be a norm closed C^* -algebra. This convention extends to three sets, thus $JJ^*J = J$ for a left ideal of a C^* -algebra as is well known. These facts follow easily from the well known results that a (norm closed) left ideal J in a C^* -algebra has a positive right contractive approximate identity. Also $J \cap J^* = J^*J \subset J \subset JJ^*$, so that J is a left ideal of JJ^* .

For the purposes of this paper we will define a *triple system* to be a norm closed subspace X of a C^* -algebra such that $XX^*X \subset X$. By ‘subspace’ we will allow for example spaces such as $B(K, H)$, regarded as the ‘1-2-corner’ of the C^* -algebra $B(H \oplus K, H \oplus K)$ in the usual way. It is well known that triple systems are ‘the same thing’ as Hilbert C^* -modules, although there is a slight difference of emphasis in the two theories; the important structure on a triple system is the ‘triple product’ xy^*z . A ‘triple subspace’ is a norm closed vector subspace of a triple system which is closed under this triple product. If X is a triple system then XX^* and X^*X are C^* -algebras, which we will call the left and right C^* -algebras of X respectively, and X is a

$(XX^*) - (X^*X)$ -bimodule. A linear map $T : X \rightarrow Y$ between triple systems is a *triple morphism* if $T(xy^*z) = T(x)T(y)^*T(z)$ for all $x, y, z \in X$. Triple systems are operator spaces, and triple morphisms are completely contractive, and indeed are completely isometric if they are 1-1 (see [13], this is related to results of Harris and Kaup). A completely isometric surjection between TRO's is a triple morphism. This last result might date back to around 1986, to Hamana and Ruan's PhD thesis independently. See [13] or [5] A.5 for a proof.

Several times in §2 we will refer to Hamana's triple envelope $\mathcal{T}(X)$ of an operator space X . This theory may be found in [13], although we review the construction of $\mathcal{T}(X)$ briefly in §1 below. An alternative account of Hamana's theory of the triple envelope is given in [5], particularly the introduction and Appendix A there. The space $\mathcal{T}(X)$ may be viewed as a triple system, and there is a canonical complete isometry $i : X \rightarrow \mathcal{T}(X)$ such that there is no nontrivial triple subspace of $\mathcal{T}(X)$ containing $i(X)$. We write $\mathcal{E}(X)$ and $\mathcal{F}(X)$ for the left- and right- C^* -algebras of $\mathcal{T}(X)$ respectively.

Next we recall the left multiplier algebra $\mathcal{M}_\ell(X)$ of an operator space X . This is a unital operator algebra, which may be viewed as a subalgebra of $CB(X)$ containing Id_X , but with a different (bigger in general) norm. Here $CB(X)$ is the space of completely bounded linear maps on X . Our first definition of $\mathcal{M}_\ell(X)$ from [5] was as the linear maps $X \rightarrow X$ corresponding to elements T in $LM(\mathcal{E}(X))$ such that $Ti(X) \subset i(X)$. The norm on $\mathcal{M}_\ell(X)$ is the $LM(\mathcal{E}(X))$ norm, and similarly for matrix norms. There are several other equivalent definitions of $\mathcal{M}_\ell(X)$ given in [5, 7, 9] - it is best to view operator space multipliers as a 'sequence of equivalent definitions'. In any particular context one or other of these definitions may be more appropriate. We will make much use the following two simple results from [6]:

Lemma 1.2. *Let A be an operator algebra with a r.c.a.i.. Then the canonical 'left regular representation' of A on itself yields completely contractive embeddings (i.e. 1-1 homomorphisms)*

$$A \hookrightarrow \mathcal{M}_\ell(A) \hookrightarrow CB(A) ,$$

and the first of these embeddings, and their composition, are completely isometric.

We remark that the canonical embedding $\mathcal{M}_\ell(A) \hookrightarrow CB(A)$, where A is an operator algebra with r.c.a.i., is not in general completely isometric, or even isometric (see Example 3.5). This has implications for our theory of multipliers in §3.

Lemma 1.3. *Suppose that $a \in B(H)$, and $\{e_\alpha\}$ is a net of contractions in $B(H)$ such that $ae_\alpha \rightarrow a$. Then $ae_\alpha e_\alpha^* \rightarrow a$, $ae_\alpha^* e_\alpha \rightarrow a$, and $ae_\alpha^* \rightarrow a$.*

2. THE IDEAL ENVELOPE

We begin by sketching Hamana's construction of the injective and the triple envelope of operator spaces. The reader who is not familiar with this will need to consult [13] for more details and notation; unfortunately the material below will be a little technical for those not versed in 'Hamana-theory'. See also the introductions of [9] and [5]). Let $X \subset B(H)$ be an operator space, and consider the Paulsen operator system

$$\mathcal{S}(X) := \begin{bmatrix} \mathbb{C}1_H & X \\ X^* & \mathbb{C}1_H \end{bmatrix} \subset M_2(B(H)) .$$

One then takes a minimal (with respect to a certain ordering) completely positive $\mathcal{S}(X)$ -projection Φ on $M_2(B(H))$. By a well-known result of Choi and Effros, $\text{Im } \Phi$ is a C^* -algebra with the multiplication $\odot$ defined by $\xi \odot \eta := \Phi(\xi\eta)$ for $\xi, \eta \in \text{Im } \Phi$, and the other algebraic operations and norm are the usual ones. One may write

$$\text{Im } \Phi = \begin{bmatrix} I_{11} & I(X) \\ I(X)^* & I_{22} \end{bmatrix} \subset M_2(B(H));$$

and $I(X)$ is an *injective envelope* of X , and I_{11}, I_{22} are injective C^* -algebras. We sometimes write I_{11} as $I_{11}(X)$, say, to emphasize the dependence on X .

We continue to think of $I_{11}, I_{22}, I(X)$ and $I(X)^*$ as subsets of $B(H)$, however the operation $\odot$ induces new products between elements of $I_{11}, I_{22}, I(X)$ and $I(X)^*$. To distinguish these multiplications from the original product on $M_2(B(H))$ we write the new products as $\circ$.

By a well known trick one may also decompose

$$\Phi = \begin{bmatrix} \psi_1 & \phi \\ \phi^* & \psi_2 \end{bmatrix}.$$

Also one may write the C^* -subalgebra of $\text{Im } \Phi$ (with the new product) generated by

$$\begin{bmatrix} 0 & X \\ 0 & 0 \end{bmatrix}$$

as

$$\begin{bmatrix} \mathcal{E}(X) & \mathcal{T}(X) \\ \mathcal{T}(X)^* & \mathcal{F}(X) \end{bmatrix} \subset \begin{bmatrix} I_{11} & I(X) \\ I(X)^* & I_{22} \end{bmatrix}.$$

This defines $\mathcal{T}(X)$, it is clearly a C^* -module or triple system (viewed as the 1-2-corner of the C^* -subalgebra just introduced, its triple product is $x \circ y^* \circ z$). Indeed the span of expressions of the form $a_1 \circ a_2^* \circ a_3 \circ a_4^* \circ \cdots \circ a_{2n+1}$, for $a_i \in X$, are dense in $\mathcal{T}(X)$. Thinking of $\mathcal{T}(X)$ as a triple system, and letting i be the canonical map $X \rightarrow \mathcal{T}(X)$, we say that $(\mathcal{T}(X), i)$ is a *triple envelope* of X .

Lemma 2.1. *Let A be a subalgebra of $B(H)$, and suppose that A has a r.c.a.i. $\{e_\alpha\}$. Then for all $a, b \in A$ we have (taking $X = A$ and using the notation above)*

$$ab = \lim_{\alpha} a \circ e_{\alpha}^* \circ b, \text{ and}$$

$$\psi_1(a) = \lim_{\alpha} a \circ e_{\alpha}^* \in I_{11}.$$

Proof. Using the notation above, and Lemma 1.3, we have

$$\psi_1(a) = \lim_{\alpha} \psi_1(ae_{\alpha}^*) = \lim_{\alpha} a \circ e_{\alpha}^*,$$

the last step by definition of the $\circ$ product. Similarly

$$ab = \phi(ab) = \phi(\lim_{\alpha} ae_{\alpha}^* b) = \lim_{\alpha} a \circ e_{\alpha}^* \circ b.$$

□

We recall that an equivalent definition of $\mathcal{M}_{\ell}(X)$ given in [9] was as $IM_{\ell}(X) = \{y \in I_{11} : y \circ X \subset X\}$. Using the last lemma we see that if A is an operator algebra with r.c.a.i., then $\psi_1(a) \in IM_{\ell}(A)$ for all $a \in A$. Thus if j is ψ_1 restricted to A , then $j : A \rightarrow IM_{\ell}(A)$. By the

last lemma, $j(a) \circ b = ab$ for all $a, b \in A$, so that j corresponds to the left regular representation $\lambda : A \rightarrow \mathcal{M}_\ell(A) \subset CB(A)$.

Proposition 2.2. *Let A be an operator algebra with r.c.a.i..*

(1) *Under the same assumptions as Lemma 2.1, and notation as above,*

$$IM_\ell(A) = \{T \in IM_\ell(A) : T \circ j(A) \subset j(A)\}.$$

(2) *Elements of $\mathcal{M}_\ell(A)$, considered as maps on A , are right A -module maps. That is, $\mathcal{M}_\ell(A) \subset CB_A(A)$ as sets. Also, $\mathcal{M}_r(A) \subset {}_A CB(A)$ as sets.*

Proof. (1) Let $a \in A$, then $T \circ j(a) = \lim_\alpha T \circ a \circ e_\alpha^*$, which equals $j(T \circ a)$ if $T \in IM_\ell(A)$. Conversely, if $T \circ j(a) = j(a')$ for an $a' \in A$, then by the last formula and 2.1 we have $T \circ j(a) \circ e_\beta = T \circ a e_\beta = a' e_\beta$, so that $T \circ a = a' \in A$.

(2) For the last assertion, if $T \in \mathcal{M}_r(A)$ then there exists $y \in I_{22}$ such that $A \circ y \subset A$, and T is the map $T(a) = a \circ y$. Hence for $c \in A$,

$$T(ca) = (ca) \circ y = \lim_\alpha c \circ e_\alpha^* \circ a \circ y = cT(a) \quad ,$$

using 2.1 twice. The other is similar. $\square$

In the statement of the following theorem, and in keeping with the usual presentation of Hamana theory (in this case the triple envelope), we forget the original product on $B(H)$ mentioned in the above discussion. That is, the products in the statement below are the $\circ$ operation in the above discussion. However in the *proof* we go back to the $\circ$ notation.

Theorem 2.3. *Let A be an operator algebra with r.c.a.i. $\{e_\alpha\}$. Suppose that $(\mathcal{T}(A), i)$ is a triple envelope for A , and let $\mathcal{E}(A) = \mathcal{T}(A)\mathcal{T}(A)^*$. The map $\psi : \mathcal{T}(A) \rightarrow \mathcal{E}(A)$ defined by $\psi(x) := \lim_\alpha x i(e_\alpha)^*$ is a well defined complete isometry, and $\psi \circ i$ is a homomorphism j on A . The range of ψ is a left ideal J of the C^* -algebra $\mathcal{E}(A)$, and ψ is also a triple morphism onto J . Thus (J, j) is another triple envelope of A . Moreover $JJ^* = \mathcal{E}(A)$, and any r.c.a.i. for A is taken by j to a r.c.a.i. for J . Finally, $\psi(x)\psi(y)^* = xy^*$ for any $x, y \in \mathcal{T}(A)$.*

Proof. By ‘abstract nonsense’ we can assume that $(\mathcal{T}(A), i)$ is the triple envelope considered above 2.1 (in short because the statement we are attempting to prove is invariant under the notion we called A -isomorphism in [5] Appendix A). We use the notation of 2.1 and above 2.1. Since elements spanning a dense subset of $\mathcal{T}(A)$ are $\circ$ products ending with a term in A , it follows from Lemma 2.1 that $\lim_\alpha x \circ e_\alpha^*$ exists in $\mathcal{E}(A)$ for any $x \in \mathcal{T}(A)$. Clearly ψ is a complete contraction extending the map j introduced above 2.2. If $x \in A$, then

$$\|\psi(x)\| = \lim_\alpha \|x \circ e_\alpha^*\| \geq \lim_\alpha \|x \circ e_\alpha^* \circ e_\beta\| = \|x e_\beta\|$$

using 2.1. Taking the limit shows that ψ is isometric on A . A similar argument using associativity of $\circ$ shows that ψ is isometric on $\mathcal{T}(A)$. Similarly, ψ is a complete isometry. If $a, b \in A$, then

$$\psi(ab) = \lim_\alpha (ab) \circ e_\alpha^* = \lim_\alpha (\lim_\beta a \circ e_\beta^* \circ b) \circ e_\alpha^* = (\lim_\beta a \circ e_\beta^*) \circ (\lim_\alpha b \circ e_\alpha^*) = \psi(a) \circ \psi(b).$$

Note also that

$$\psi(a) \circ \psi(b)^* = \lim_\alpha \lim_\beta a \circ e_\alpha^* \circ e_\beta \circ b^* = \lim_\beta (a e_\beta) \circ b^* = a \circ b^* \quad .$$

Thus in the language of the statement of the theorem $\psi(i(a))\psi(i(b))^* = i(a)i(b)^*$.

By looking at the natural dense subsets of $\mathcal{T}(A)$ and $\mathcal{E}(A)$ it is easy to argue that the range of ψ is a left ideal J . By a fact mentioned in the introduction this implies that ψ is a triple morphism. Since j is the restriction of ψ to A , the remaining assertions are easy. One needs to use the fact that ψ is a triple morphism, and that $\mathcal{T}(A)$ and therefore consequently J , have dense subsets spanned by terms which are alternating products as mentioned above 2.1. $\square$

Theorem 2.4. *Let A be an operator algebra with a r.c.a.i., let $(I(A), i)$ be its injective envelope discussed above 2.1 say, and let B be the injective unital C^* -algebra I_{11} (also discussed above). Then there is an orthogonal projection $e \in B$, such that $I'(A) = (Be, j)$ is also an injective envelope for A , where j is as above. For this new injective envelope, $I'_{22} \cong eBe$ and $I'_{11} \cong B$. The completely isometric embedding $j : A \rightarrow I'(A) \subset B$ is a homomorphism. Moreover if A has a right identity f then $j(f) = e$.*

Before we prove this, we remark that we may define the *ideal injective envelope* of an operator algebra A with r.c.a.i., to be $I'(A)$ as in the theorem.

Proof. Define $\psi : I_{11} \circ A \rightarrow I_{11}$ by $\psi(x) = \lim_{\alpha} x \circ e_{\alpha}^*$. By 2.1 this limit exists and ψ is clearly completely contractive. Note that ψ is also a left I_{11} -module map, so that by a result in [9] we may extend ψ to a completely contractive left I_{11} -module map $\hat{\psi} : I(A) \rightarrow I_{11}$. The restriction of this map to A was the map called j above. By Hamana's 'essential property' for $I(A)$ [13], $\hat{\psi}$ is a complete isometry. By [9] 2.7, there is a $v \in \text{Ball}(I(A))$ such that $\hat{\psi}(x) = x \circ v^*$, for all $x \in I(A)$. Also its range is a left ideal of I_{11} , so that $\hat{\psi}$ is a triple morphism by a fact at the end of the introduction.

Next define a map $I_{11} \circ j(A) \subset I_{11} \rightarrow I(A)$ by $z \mapsto \lim_{\alpha} z \circ e_{\alpha}$. To see that this limit exists note that for $x \in I_{11}$ and $a \in A$ we have by the type of calculations found in the proof of 2.3, that $\lim_{\alpha} x \circ j(a) \circ e_{\alpha} = x \circ a$. So the limit exists on $I_{11} \circ j(A)$, and moreover this map is exactly the map $x \circ j(a) \mapsto x \circ a$. It is thus a completely contractive left I_{11} -module map, and extends by [9] to a completely contractive left I_{11} -module map $\mu : I_{11} \rightarrow I(A)$. There clearly exists $w \in \text{Ball}(I(A))$ such that $\mu = - \circ w$. Now notice that the composition $\mu\hat{\psi}$ is a completely contractive map $I(A) \rightarrow I(A)$, which restricts to the identity map on A . By Hamana's rigidity property [13], $\mu\hat{\psi} = \text{Id}$ on $I(A)$. Thus $\hat{\psi}$ is a complete isometry, μ is onto. Moreover since $x \circ (v^* \circ w) = x$ for all $x \in I(X)$, it follows from [9] Corollary 1.3, that $v^* \circ w$ is the identity of I_{22} . Thus by a well know operator theory fact, $v = w$. We define $e = v \circ v^* \in I_{11} = B$, this is an orthogonal projection. Moreover $\text{Ran } \hat{\psi} = Be$, as may be seen easily from the above. It follows immediately that (Be, j) is an injective envelope for A , where $j(a) = a \circ v^*$ (which coincides with the map j of previous results), which is a homomorphism.

Note that by [9] Theorem 1.8, $I'_{11} \cong M(I'(X)I'(X)^*) = M(I(X)I(X)^*) \cong I_{11}$, and $I'_{22} \cong M(eBe) = eBe$.

Finally, if A has right identity f then we may take v above to be f , so that $j(f) = f \circ f = e$. $\square$

It is clear from 2.3 that for an operator algebra A with r.c.a.i., $\mathcal{T}(A)$ may be taken to be a left ideal in a C^* -algebra, and this left ideal therefore possesses Hamana's universal property of the

triple envelope (see [13] or [5] Appendix A). However examining this property in this case gives an interesting refinement, by viewing $(\mathcal{T}(A), j)$ as a *left ideal extension* of A .

To be more specific we say that a pair (J, i) consisting of a left ideal J in a C^* -algebra, and a completely isometric homomorphism $i : A \rightarrow J$, is a *left ideal extension* of A if $i(A)$ ‘generates J as a triple system’. That is, the span of expressions of the form $i(a_1)i(a_2)^*i(a_3)i(a_4)^*\cdots i(a_{2n+1})$, for $a_i \in A$, are dense in J . It follows from this that $\{i(e_\alpha)\}$ is a r.c.a.i. for J if $\{e_\alpha\}$ is a r.c.a.i. for A . Then it is clear from 2.3 that $(\mathcal{T}(A), j)$ may be taken to be a left ideal extension of A , and it is the ‘minimal such’. The new point in the theorem below is that τ may be chosen to be a homomorphism:

Corollary 2.5. *Let A be an operator algebra with r.c.a.i.. Then there exists a left ideal extension $(\mathfrak{J}_e(A), j)$ of A , with $\mathfrak{J}_e(A)$ a left ideal in a C^* -algebra $\mathcal{E}(A) = \mathfrak{J}_e(A)\mathfrak{J}_e(A)^*$, such that for any other left ideal extension (J, i) of A , there exists a (necessarily unique and surjective) completely contractive homomorphism $\tau : J \rightarrow \mathfrak{J}_e(A)$, which is also a triple morphism, such that $\tau \circ i = j$. Thus $\mathfrak{J}_e(A)/(\ker \tau) \cong J$ completely isometrically homomorphically (i.e as operator algebras) too. Moreover $(\mathfrak{J}_e(A), j)$ is unique in the following sense: given any other (J', j') with this universal property, then there exists a surjective completely isometric homomorphism $\theta : \mathfrak{J}_e(A) \rightarrow J'$ such that $\theta \circ j = j'$.*

Proof. The uniqueness is fairly obvious and standard. The existence follows from 2.3 (setting $\mathfrak{J}_e(A) = (J, j)$ there), together with Hamana’s universal property for the ‘triple envelope’ mentioned above. Indeed we have:

$$\tau(xy) = \lim \tau(xi(e_\alpha)^*y) = \lim \tau(x)j(e_\alpha)^*\tau(y) = \tau(x)\tau(y)$$

for $x, y \in J$, using Lemma 1.3 and the observations above. □

We call $(\mathfrak{J}_e(A), j)$ the *left ideal envelope* of A , and continue to write $\mathcal{E}(A) = \mathfrak{J}_e(A)\mathfrak{J}_e(A)^*$. Again, j will be called the *Shilov embedding homomorphism*.

Corollary 2.6. *Let A be an operator algebra with r.c.a.i., and let $j : A \rightarrow \mathcal{E}(A)$ be the Shilov embedding homomorphism. Then $\mathcal{M}_\ell(A) \cong \{T \in LM(\mathcal{E}(A)) : Tj(A) \subset j(A)\}$ completely isometrically isomorphically.*

This last corollary above is a restatement of 2.2, using the definition of $\mathcal{M}_\ell(A)$ given in the introduction.

3. THE LEFT MULTIPLIER ALGEBRA OF AN ALGEBRA WITH RIGHT APPROXIMATE IDENTITY

The readers first thought here might be that this case must be analogous to the case of the left multiplier algebra of an operator algebra with l.c.a.i., which was treated in [6]. However in fact the theory is rather different. Part of this may be seen by considering the basic example of $A = C_n$ (the $n \times n$ matrices ‘supported on’ the first column). It is fairly clear that its left multiplier operator algebra ‘should be’ M_n .

After a little thought about operator algebras with r.c.a.i., one comes up with the following list (which we shall refer to throughout this section) of ‘possible candidate definitions’ for $LM(A)$ in this case (all of which contain A completely isometrically):

- (I) $\{x \in A^{**} : x\hat{A} \subset \hat{A}\}$,
- (II) $\{T \in B(H) : T\pi(A) \subset \pi(A)\}$, where $\pi : A \rightarrow B(H)$ is a completely isometric nondegenerate representation,
- (III) $\mathcal{M}_\ell(A)$,
- (IV) $CB_A(A)$ (right module maps),
- (V) the ‘maximum essential left multiplier extension of A ’.

We will spell out later what is meant in this setting by ‘essential left multiplier extension’. One can rule out (I) fairly quickly as a plausible candidate since it is not unital (e.g. take $A = C_n$). However one would hope that most of the other four items are completely isometrically isomorphic. Unfortunately, most of the equivalences amongst these items break down if A is a general operator algebra with r.c.a.i.. Nonetheless one of these five candidates will emerge from our study below as the ‘winner’, namely as the appropriate version of $LM(A)$ in this case. This seems quite interesting.

In the special case of a left ideal J in a C^* -algebra things are much better. Most of these five items are then completely isometrically isomorphic. Indeed in this case we shall see that most of these coincide with the usual left multiplier algebra $LM(JJ^*)$ of the C^* -algebra JJ^* . One may view this as being predicted by the theory of Hilbert C^* -modules. See also [14]. Indeed the analogous ‘left multiplier C^* -algebra’ $LM^*(J)$, which may be defined to be $M(JJ^*)$, or equivalently $A_l(J)$ in the notation of [5], also has a very satisfactory theory which has essentially been done in [14]. Hence we shall not mention $LM^*(J)$ again here, leaving any details of its theory to the interested reader.

In this special case of a left ideal J in a C^* -algebra we will take the term ‘essential left multiplier extension’ used in (V), to be defined as in the proof of (4) in the main theorem in [6] §4. Of particular interest, in (II), are the *ideal representations* of J discussed in [6], that is, the completely isometric representations $\pi : J \rightarrow B(H)$ which are restrictions of faithful $*$ -representations $JJ^* \rightarrow B(H)$ (or equivalently, the completely isometric representations of J which are also triple morphisms).

Theorem 3.1. *Let J be a left ideal of a C^* -algebra. The algebras in (III)-(V) are all completely isometrically isomorphic to $LM(JJ^*)$. Any algebra in (II)-(V) contain the algebra in (I) completely isometrically isomorphically as a (proper, in general) subalgebra. If, further, in (II) we only consider 1-1 nondegenerate ideal representations π , then the algebras in (II)-(V) are all completely isometrically isomorphic. Also $B_J(J) = CB_J(J)$ isometrically.*

Proof. In this case J is a C^* -module, and $\mathcal{M}_\ell(J)$ may easily be seen, from [5] A.4 if necessary, to be $LM(JJ^*)$. Clearly $CB_J(J) \subset CB_{J^*J}(J)$, and conversely, if $T \in CB_{J^*J}(J)$ then for $x, y \in J$ we have using Lemma 1.3 that $T(xy) = \lim T(xe_\alpha^*y) = \lim T(x)e_\alpha^*y = T(x)y$, if $\{e_\alpha\}$ is the r.c.a.i. in J . Thus $CB_J(J) = CB_{J^*J}(J)$. By a theorem of Lin ([15] 1.4) $B_{J^*J}(J) = LM(JJ^*)$, and the operator space version of this is true too (see [3]), that is $CB_{J^*J}(J) = LM(JJ^*)$ completely isometrically. Thus (III) = (IV), and we obtain the last assertion of the statement of the theorem too. Looking at the definition of a ‘essential left multiplier extension’, and the associated ‘ordering’, below, or defining these as we did in the proof of (4) in the main theorem in [6] §4, it is now clear that (IV) is an essential left multiplier extension, so that it is the biggest such. That is, (IV) = (V).

We will defer until 3.2 the proof that an algebra in (II) contains the algebra in (I) completely isometrically. Finally, given a nondegenerate faithful $*$ -representation $\theta : JJ^* \rightarrow B(H)$, we know from the multiplier theory for C^* -algebras (see e.g. [19] 3.1.12) that $LM(JJ^*) = \{T \in B(H) : T\theta(JJ^*) \subset \theta(JJ^*)\}$. However since for any ideal J we have $J = JJ^*J$, if $T \in B(H)$, then $T\theta(J) \subset \theta(J)$ if and only if $T\theta(JJ^*) \subset \theta(JJ^*)$. Thus the algebra in (II) equals $LM(JJ^*)$ too in this case. $\square$

We have now completed our discussion of $LM(J)$ for a left ideal J in a C^* -algebra, and turn to the more general case of an arbitrary operator algebra A with r.c.a.i.. Since this case is the most complicated, we will be a little more formal and rigorous in our presentation below.

We begin by defining a *left multiplier extension* of A to be a pair (B, π) consisting of an operator algebra B with an identity of norm 1, and a completely isometric homomorphism $\pi : A \rightarrow B$ such that $\pi(A)$ is a left ideal of B . We say that (B, π) is an *essential left multiplier extension* of A if in addition the canonical map $B \rightarrow CB_A(A)$ is 1-1. Note that this canonical map is then a completely contractive homomorphism into $CB_A(A)$ (viewing A as a right A -module). In the 2-sided case $CB_A(A)$ is itself a left multiplier extension of A , and is therefore the largest left multiplier extension, but this is not true in general for us (see example 3.4). We next define an ordering on essential left multiplier extensions of A : namely $(B, \pi) \leq (B', \pi')$ if there exists a (necessarily unique and one-to-one) completely contractive homomorphism $\theta : B \rightarrow B'$ such that $\theta \circ \pi = \pi'$. We remark that in the theory in [6] §4 or for left ideals in C^* -algebras one may insist that θ is completely isometric, but we are not certain if we can make this requirement in general. Note that this ordering says that B may be viewed as a subalgebra of B' , but with a possibly bigger norm.

We will use the fact that an essential left multiplier extension (B, π) has the following rigidity property: if $\theta : B \rightarrow B$ is a completely contractive homomorphism extending the identity mapping on $\pi(A)$, then θ is the identity mapping (To see this note that $\theta(b)\pi(a) = \theta(b\pi(a)) = b\pi(a)$, so that $\theta(b) - b$ is in the kernel of the canonical map $B \rightarrow CB(A)$).

We say that two left multiplier extensions (B, π) and (B', π') are *A-equivalent* if there exists a unital completely isometric isomorphism $\theta : B \rightarrow B'$ with $\theta \circ \pi = \pi'$. Note that this is an equivalence relation, and that ' $\leq$ ' induces a well defined ordering on the equivalence classes. It follows that if there exists a maximum essential left multiplier extension of A , then it is unique up to A -equivalence. We remark further that if two left multiplier extensions are A -equivalent, and if one is essential, then so is the other. We leave the proof of this as an easy exercise.

One more definition: we say that a representation $\pi : A \rightarrow B(H)$ is a completely isometric *Shilov representation*, if π is the restriction of a 1-1 $*$ -homomorphism $\mathcal{E}(A) \rightarrow B(H)$ (viewing $A \subset \mathcal{E}(A)$ as in 2.5 say via the completely isometric homomorphism j there). Note that the 'ideal representations' π considered in Theorem 3.1 happen also to be Shilov representations.

Theorem 3.2. *Let A be an operator algebra with r.c.a.i.. Then $\mathcal{M}_\ell(A)$ is the (unique up to A -equivalence) maximum essential left multiplier extension of A . That is, the algebras in items (III) and (V) in the list at the beginning of §5 are completely isometrically isomorphic. Also $\mathcal{M}_\ell(A)$ contains, and dominates in the ordering above, the essential left multiplier extensions described in (II), which in turn contain the algebra in (I) completely isometrically isomorphically. However if π in (II) is a Shilov representation of A then the algebra in (II) is completely isometrically isomorphic to (III) and (V). The algebra in (IV), namely $CB_A(A)$, is not in general an operator*

algebra, and its subalgebra which corresponds to the algebra in (III) (and (V)) does not correspond isometrically. That is, the canonical 1-1 homomorphic embedding of $\mathcal{M}_\ell(A)$ into $CB_A(A)$ (or even $B_A(A)$) need not be isometric.

Proof. That the canonical inclusion of (III) in (IV) is not isometric may be seen in Example 3.5. That $CB_A(A)$ is not in general an operator algebra may be seen in Example 3.4.

We have seen in 2.2 that $(\mathcal{M}_\ell(A), \lambda)$ is an essential left multiplier extension. It is evident from the fundamental properties of $\mathcal{M}_\ell(A)$ (from Theorem 4.10 (1) of [5] say) that given any essential left multiplier extension (B, π) of A , there is a canonical completely contractive homomorphism $\sigma : B \rightarrow \mathcal{M}_\ell(A)$. It is obvious that via this homomorphism $(B, \pi) \leq (\mathcal{M}_\ell(A), \lambda)$. So $(\mathcal{M}_\ell(A), \lambda)$ is the maximum essential left multiplier extension.

If $\pi : A \rightarrow B(H)$ is a completely isometric ‘nondegenerate representation’ (by which we mean at least that $\pi(A)H$ is dense in H), then $\{T \in B(H) : T\pi(A) \subset \pi(A)\}$ is fairly evidently an essential left multiplier extension of A , and hence is dominated by $\mathcal{M}_\ell(A)$ (by the fact in the previous paragraph).

We now show any algebra in (II) contains (I) completely isometrically. If $\pi : A \rightarrow B(H)$ is any (not necessarily nondegenerate) completely isometric homomorphism, consider the following sequence of completely contractive homomorphisms:

$$A \hookrightarrow A^{**} \xrightarrow{\pi^{**}} B(H)^{**} \rightarrow B(H)$$

where the first and last maps here are the canonical inclusion and projection. Let σ be the composition of the last two maps, restricted to $\{x \in A^{**} : x\hat{A} \subset \hat{A}\}$. Then $\sigma(\hat{a}) = \pi(a)$ for $a \in A$, and so for x as above and $a \in A$ we have

$$\sigma(x)\pi(a) = \sigma(x\hat{a}) \in \sigma(\hat{A}) \subset \pi(A).$$

Thus σ maps into the left idealizer of $\pi(A)$ in $B(H)$. To see that σ is isometric note that

$$\|\sigma(x)\| \geq \|\sigma(x)\pi(e_\alpha)\| = \|\sigma(x\hat{e}_\alpha)\| = \|\pi(x\hat{e}_\alpha)\| = \|x\hat{e}_\alpha\|.$$

Now 1 is a weak* limit point of the $\{\hat{e}_\alpha\}$, and using the separate weak* continuity of the product on A^{**} , we see that $\|\sigma(x)\| \geq \|x\|$. Similarly σ is completely isometric.

Finally we discuss the equivalence of (III) with (II) for Shilov representations π . Take $\pi : \mathcal{E}(A) \rightarrow B(H)$ to be any nondegenerate 1-1 *-homomorphism, then it is easy to see that $\pi|_A$ is a nondegenerate completely isometric homomorphism. By the basic theory of multipliers of C^* -algebras we may view $LM(\mathcal{E}(A)) = \{T \in B(H) : T\pi(\mathcal{E}(A)) \subset \pi(\mathcal{E}(A))\}$. From 2.6 we thus have $\mathcal{M}_\ell(A)$ completely isometrically isomorphic to

$$\{T \in B(H) : T\pi(\mathcal{E}(A)) \subset \pi(\mathcal{E}(A)), \text{ and } T\pi(A) \subset \pi(A)\} = \{T \in B(H) : T\pi(A) \subset \pi(A)\}$$

since $j(A)j(A)^*$ generates $\mathcal{E}(A)$ as a C^* -algebra. □

Proposition 3.3. *Let A be an operator algebra with r.c.a.i.. Then $CB_A(A)$ is an operator algebra if and only if $CB_A(A)$ is completely isometrically isomorphic to $\mathcal{M}_\ell(A)$.*

Proof. The one direction is trivial since $\mathcal{M}_\ell(A)$ is an operator algebra. To see the other note that by 1.2 and 2.2 we have $\lambda(A) \subset CB_A(A)$, and $\mathcal{M}_\ell(A) \subset CB_A(A)$ although the $\mathcal{M}_\ell(A)$ -norm is possibly larger than the $CB_A(A)$ -norm. Thus if $CB_A(A)$ is an operator algebra then

$(CB_A(A), \lambda)$ is an essential left multiplier extension of A a 1-1 completely contractive homomorphism $CB_A(A) \rightarrow \mathcal{M}_\ell(A)$ (by the equivalence of (II) and (V) in 3.2). It is easy to see that this homomorphism restricts to the identity map on $\mathcal{M}_\ell(A)$, whose inverse is completely contractive as a map $\mathcal{M}_\ell(A) \rightarrow CB_A(A)$. $\square$

Next we consider some examples:

Example 3.4.

Let $A \subset M_3$ be the subalgebra supported on the 1-2, 1-3, 2-2, and 3-3 entries only:

$$A = \begin{bmatrix} 0 & * & * \\ 0 & * & 0 \\ 0 & 0 & * \end{bmatrix}.$$

Then A is an operator algebra with right identity of norm 1. One may easily compute $\mathfrak{J}_e(A) = \mathcal{T}(A) = I(A) = M_{3,2}$, and $\mathcal{E}(A) = M_3$, from which it follows that $\mathcal{M}_\ell(A)$ is the subalgebra of M_3 spanned by A and I_3 . Thus $\mathcal{M}_\ell(A)$ is 5 dimensional. Indeed in this example, $\mathcal{M}_\ell(A)$ coincides with what we called the ‘multiplier unitization’ of A . On the other hand the algebra given by item (I) in the list at the beginning of §5, is A again, which is 4 dimensional. A tedious but completely elementary algebraic computation shows that $CB_A(A)$, item (IV) on the aforementioned list, is 8 dimensional. This shows by 3.3 that $CB_A(A)$ cannot be completely isometrically isomorphic to an operator algebra (in contrast to the 2-sided c.a.i. case).

Example 3.5.

It is perhaps true in the previous example that $\mathcal{M}_\ell(A)$ is not contained isometrically in $CB(A)$, but this seems difficult to check. Instead we generalize this example to an interesting example of an operator algebra A with right identity of norm 1, for which $\mathcal{M}_\ell(A)$ is fairly clearly not contained isometrically in $CB(A)$ (or $B(A)$). To do this we take a C^* -algebra B with no nontrivial 2-sided ideals, and a closed subspace $X \subset B$ which generates B as a triple system. To be more concrete one could take $B = M_3$ and $X \subset M_3$ as in Example 4.4 of [5]. Consider $A \subset M_3(B)$ the subalgebra supported on the 1-2, 1-3, 2-2, and 3-3 entries only (i.e. zero elsewhere), which has scalar multiples of the identity operator in the 2-2 and 3-3 entries, and has operators from X in the 1-2 and 1-3 entries. If one computes the triple system generated by A in $M_3(B)$, one sees that one obtains $M_{3,2}(B)$ as in the previous example. As in that example one sees that $\mathcal{E}(A) = M_3(B)$ and $\mathcal{M}_\ell(A)$ is the subalgebra of $M_3(B)$ spanned by A and matrices supported only in the 1-1 entry, and that entry may be anything in $\mathcal{M}_\ell(X)$. Now consider one of these last matrices, T say, and let a be its one nonzero entry (in $\mathcal{M}_\ell(X)$). Viewing $T \in CB(A)$ as a left multiplication map, it is clear that its norm or ‘cb-norm’ is the same as the norm or ‘cb-norm’ of its 1-1 entry, as a left multiplication map on X . But in Example 4.4 of [5] we showed that neither of these numbers equals $\|a\| = \|T\|$ in general.

Remark. If A is an operator algebra with r.c.a.i., then there is a most important element of A^{**} , namely the right identity E there. A very natural question arises as to whether $E \in LM(A)$, and more particularly whether $E\hat{A} \subset \hat{A}$? In fact this is not true in general. To see this suppose that we have an operator algebra A with r.c.a.i. but no l.c.a.i., which has property $(\mathcal{R})$ discussed in [6]. If $E\hat{A} \subset \hat{A}$ then in the notation of 2.2 there is a corresponding element of $e \in IM_l(A)$

such that $e \circ a = Ea$ for $a \in A$. In particular for b in the algebra $\mathcal{R}(A)$ defined in [6] §2, we have $e \circ b = b$. If one chooses an A with $I(\mathcal{R}(A)) = I(A)$ (such examples exist) then by the ‘rigidity property’ of the injective envelope [13], it follows that $e \circ a = a$ for all $a \in I(A)$. Thus in particular, for $a \in A$ we have $Ea = a$, from which it follows by [10] 28.7 that a has a l.c.a.i., which is a contradiction.

Remark. (Unitizations.) Notice that any time one has a (essential) left multiplier extension (B, i) of an operator algebra A with r.c.a.i., we get a unitization of A defined as $i(A) + \mathbb{C}1_B$. Conversely, any unitization of A should be a left multiplier extension of A .

Taking the multiplier extension $\mathcal{M}_\ell(A)$ here, let us call the corresponding unitization the ‘Shilov unitization’ A^1 of A ; it will be the unitization with the smallest norm. Of course another universal essential left multiplier extension may be constructed by considering a left idealizer in the direct sum of all essential left multiplier extensions of A . We obtain an associated unitization of A , this one with the largest norm. Indeed this may also be described by simply assigning A^1 the supremum of all operator algebra structures on A^1 coinciding with the usual one on A .

We do not know in general if there is a simple formula for the norm on the unitization in terms of the norm on A , as there is in the case of an algebra with two-sided c.a.i.. Of course if the span of Id_X and $\lambda(A)$ inside $CB_A(A)$ is an operator algebra, then this coincides with the ‘Shilov unitization’ mentioned above, by an argument similar to 3.3. These matters deserve investigation.

Finally, we end with some remarks on the 2-sided multiplier algebra $M(A)$ of an operator algebra with l.c.a.i. say. One natural candidate for $M(A)$ might be constructed as follows: take any candidate (D, μ) for $LM(A)$ (as in [6] §4), and define $M(A) = \{x \in D : \mu(A)x \subset \mu(A)\}$. It is easy to see that if we take a (D', μ') which is A -isomorphic to (D, μ) , then one obtains the same $M(A)$, up to completely isometric isomorphism, or up to a suitable notion of A -equivalence). Perhaps this coincides with $M(\mathcal{L}(A))$ if A has property $(\mathcal{L})$.

One would hope that one might get the same algebra $M(A)$ by looking at $\{x \in M_r(A) : x\rho(A) \subset \rho(A)\}$ where ρ is the ‘right regular antirepresentation’ of A on itself, but this is not correct (consider R_2). In fact this algebra is related to a ‘big’ multiplier algebra $BM(A)$ of A , defined as the ‘double idealizer’ of $j(A)$ in $M(B)$, where B is the injective C^* -algebra in 2.4. Another approach to $M(A)$ is via ‘double centralizers’ as in [21] and [16] §1.2. There must be some relations between all these candidates for $M(A)$, and some of the algebras discussed at the end of [6] §4 will also play a role. The second author is currently investigating these matters to see if there are any satisfactory results here.

Acknowledgement: Several results in this paper will be included as part of the second author’s Ph. D. thesis, done under the direction of the first author during the ’00-’01 academic year. He expresses his gratitude to the first author for his kind instruction and advising.

REFERENCES

- [1] W. B. Arveson, *Subalgebras of C^* -algebras*, Acta Math. **123** (1969), 141-224.
- [2] D. P. Blecher, *Some general theory of operator algebras and their modules*, in *Operator algebras and applications*, A. Katavolos (editor), NATO ASIC, Vol. 495, Kluwer, Dordrecht, 1997.

- [3] D. P. Blecher, *A new approach to C^* -modules*, Math. Annalen **307** (1997), 253-290.
- [4] D. P. Blecher, *Unpublished notes on multiplier operator algebras*, (2000).
- [5] D. P. Blecher, *The Shilov boundary of an operator space, and the characterization theorems*, J. Funct. An. **182** (2001), 280-343.
- [6] D. P. Blecher, *One-sided ideals and approximate identities in operator algebras*, Preprint (September '01).
- [7] D. P. Blecher, E. G. Effros and V. Zarikian, *One-sided M -ideals and multipliers in operator spaces I*, Preprint (2000). To appear *Pacific J. Math.*
- [8] D. P. Blecher, P. S. Muhly and V. I. Paulsen, *Categories of operator modules - Morita equivalence and projective modules*, *Memoirs of the American Mathematical Society* Vol. 143, number 681 (Jan, 2000).
- [9] D. P. Blecher and V. I. Paulsen, *Multipliers of operator spaces, and the injective envelope*, Pacific J. Math. **200** (2001), 1-17.
- [10] F. F. Bonsall and J. Duncan, *Complete normed algebras*, Springer-Verlag, New York-Heidelberg (1973).
- [11] E. G. Effros and Z. J. Ruan, *Operator Spaces*, Oxford University Press (2000).
- [12] M. Hamana, *Injective envelopes of operator systems*, Publ. R.I.M.S. Kyoto Univ. **15** (1979), 773-785.
- [13] M. Hamana, *Triple envelopes and Silov boundaries of operator spaces*, Math. J. Toyama University **22** (1999), 77-93.
- [14] M. Kaneda and V. I. Paulsen, *Characterizations of essential ideals as operator modules over C^* -algebras*, Preprint (2000).
- [15] H. Lin, *Bounded module maps and pure completely positive maps*, J. Operator Theory **26** (1991), 121-138.
- [16] T. W. Palmer, *Banach Algebras and the general theory of $*$ -algebras Vol. I*, Encyclopedia of mathematics and its applications Vol. 49, Cambridge University Press (1994).
- [17] V. I. Paulsen, *Completely bounded maps and dilations*, Pitman Research Notes in Math., Longman, London, 1986.
- [18] V. I. Paulsen, *Unpublished notes on multiplier algebras*, (circa 1992).
- [19] G. Pederson, *C^* -algebras and their automorphism groups*, Academic Press (1979).
- [20] G. Pisier, *Introduction to the theory of operator spaces*, to appear.
- [21] Y-t. Poon and Z. J. Ruan, *Operator algebras with contractive approximate identities*, Canadian J. Math. **46**(1994), 397-414.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HOUSTON, 4800 CALHOUN, HOUSTON, TX 77204-3008, U.S.A.

E-mail address: dblecher@math.uh.edu and kaneda@math.uh.edu